

Proceedings

Open Access

Decreased Capacity for Vitamin A Storage in Hepatic Stellate Cells for Arctic Animals

Haruki Senoo*¹, Kenjiro Wake², Heidi L Wold³, Nobuyo Higashi¹, Katsuyuki Imai¹, Jan Øivind Moskaug³, Naosuke Kojima¹, Mitsutaka Miura¹, Takeya Sato¹, Mitsuru Sato¹, Norbert Roos⁴, Trond Berg⁴, Kaare R Norum³ and Rune Blomhoff³

Address: ¹Department of Anatomy, Akita University School of Medicine, Akita, 010-8543, Japan, ²Liver Research Unit, Minophagen Pharmaceutical Co. Ltd., Tokyo, 160-0004, Japan, ³Institute for Nutrition Research, Faculty of Medicine, University of Oslo, Oslo, Norway and ⁴Department of Biology, University of Oslo, Oslo, Norway

Email: Haruki Senoo* - senoo@ipc.akita-u.ac.jp; Kenjiro Wake - wake@minophagen.co.jp; Heidi L Wold - heidilwold@hotmail.com; Nobuyo Higashi - nobuyo@med.akita-u.ac.jp; Katsuyuki Imai - imai@med.akita-u.ac.jp; Jan Øivind Moskaug - j.o.moskaug@basalmed.uio.no; Naosuke Kojima - nkojima@med.akita-u.ac.jp; Mitsutaka Miura - mttk@med.akita-u.ac.jp; Takeya Sato - tksato@med.akita-u.ac.jp; Mitsuru Sato - mitsuru@med.akita-u.ac.jp; Norbert Roos - norbert.roos@bio.uio.no; Trond Berg - trond.berg@bio.uio.no; Kaare R Norum - k.r.norum@basalmed.uio.no; Rune Blomhoff - rune.blomhoff@basalmed.uio.no

* Corresponding author

from 11th International Symposium on the Cells of the Hepatic Sinusoid and their Relation to Other Cells
Tucson, Arizona, USA, 25–29 August, 2002

Published: 14 January 2004

Comparative Hepatology 2004, **3**(Suppl 1):S18

This article is available from: <http://www.comparative-hepatology.com/content/3/S1/S18>

Introduction

Under physiological conditions, hepatic stellate cells store 80 % of the total vitamin A in the whole body as retinyl palmitate in lipid droplets in the cytoplasm, and regulate both transport and storage of vitamin A [1-3].

It has been demonstrated that animal or human individuals exposed to drugs such as methadone, prednisone and phenobarbital, antiepileptics [4], and xenobiotics like the environmental contaminants DDT, PCD and dioxins have dramatic changes in their retinoid metabolism and function.

Recent data demonstrate that top predators among Svalbard mammals and birds like polar bear, arctic fox and glaucous gull accumulate relatively large amounts of persistent organic pollutants [5]. Since it has been reported that vitamin A accumulates at near toxic doses in some arctic predators [6] and recent data suggest that PCB and DDT may reduce the threshold for vitamin A-toxicity [7], an increasing accumulation of persistent organic pollutants might eventually precipitate vitamin A-toxicity in these animals. To elucidate the possibility of vitamin A-related toxicity in arctic predators, we have performed a

systematic characterization of the hepatic vitamin A-storage, which is the best index of the vitamin A-status, in mammals of the Svalbard archipelago.

Methods

After getting permission to hunt the animals from the district governor of Svalbard, 11 arctic foxes and 14 bearded seals, were caught in the Svalbard archipelago near Longyearbyen (78° N, 15° E) in the period from August 1996 to September 2001. Three polar bears were shot in self-defense at Svalbard February and August 1998 in Ny Ålesund and Hornsund. Distribution and content of vitamin A in livers and other organs were analyzed by morphological methods such as transmission electron microscopy, fluorescence microscopy for detection of autofluorescence of vitamin A and gold chloride staining [8] and high-performance liquid chromatography.

Results and Discussion

The amounts of vitamin A stored in livers of arctic animal are shown in Table 1. The median values are presented due to the relatively large individual variations. These values are much higher than all other arctic animals studied as well as their genetically related continental top

Table 1: Total retinol concentration in liver and kidney from rats and arctic animals.

Species	Total retinol concentration ^a		
	Liver (nmol/g)	Kidney (nmol/g)	Kidney concentration as percent of liver concentration
Control rats (n = 3) ^b	1106	9	0.81
Vitamin A-fed rats (n = 3) ^b	16877	224	1.33
Arctic animals:			
Polar bear (n = 3)	18279	102	0.56
Arctic fox (n = 8)	18641	1623	8.71
Bearded seal (n = 10)	4652	39	0.84

^a Total retinol content (i.e. the sum of retinol and retinyl esters) is presented as mean for rats and median for arctic animals. ^b These results have been published in a previous study [9].

predators (data not shown). The values are also high compared to normal human values and experimental animals like mouse and rat (200–600 nmol per gram) [9].

The concentration of total retinol in kidney is normally less than 1% of the concentration in liver. However, following an intake of excessive amounts of vitamin A or after experimental treatment with xenobiotics like HCB, TCDD and PCB, the concentration in kidneys may increase several-fold [10,11]. Kidney total retinol may therefore be used as a biomarker for vitamin A-related toxicity or excess. When we measured the kidney concentration of total retinol in the top arctic predators we observed that polar bear and bearded seal had kidney levels below 1% of their liver value (Table 1). Arctic fox, however, had kidney levels of about 9% of the liver values (Table 1).

Strong autofluorescence and gold chloride staining were present in hepatic stellate cells of polar bear, arctic fox, and bearded seal (data not shown). The distribution of stored vitamin A in the arctic animals was essentially the same as that published previously in normal rat and human liver [2,3]. In livers of arctic fox, polar bear, and bearded seal, stellate cells stored one or two large lipid droplets in their cytoplasm (data not shown). In the kidney of arctic foxes and bearded seals, interstitial cells (renal stellate cells) stored vitamin A-lipid droplets (data not shown).

Increased kidney concentrations of total retinol in arctic fox most likely are a sign of pollutant-induced vitamin A-toxicity. It is interesting to note that the highest accumulation of organic pollutants in arctic animals occur in the arctic fox [12]. Relative decrease of liver concentration of total retinol in arctic fox might be due to the decreased capacity for storage of vitamin A in hepatic stellate cells.

Acknowledgements

The authors are grateful to Jørn Eldar Fortun, Anders Friberg, and Trond Østaa for their excellent hunting. We also thank the University Courses at Svalbard for their generous assistance by allowing us to use their laboratory facilities in Longyearbyen, the Norwegian Polar Institute and the District Governor at Svalbard.

References

- Blomhoff R: **Vitamin A in Health and Disease**. New York, Marcel Dekker Inc 1994.
- Wake K: **Sternzellen in the liver, perisinusoidal cells with special reference to storage of vitamin A**. *Am J Anat* 1971, **132**:429-462.
- Blomhoff R, Wake K: **Perisinusoidal stellate cells of the liver: important roles in retinol metabolism and fibrosis**. *FASEB J* 1991, **5**:271-277.
- Botilsrud M, Wathne KO, Norum KR, Blomhoff R: **Effects of anti-epileptics on the hepatic storage of retinol**. *In Vivo* 1988, **2**:253-256.
- Muir DC, Norstrom RJ: **Geographical differences and time trends of persistent organic pollutants in the Arctic**. *Toxicol Lett* 2000, **112-113**:93-101.
- Rodahl K: **Toxicity of polar bear liver**. *Nature* 1949, **164**:530-531.
- Norstrom RJ, Muir DC: **Chlorinated hydrocarbon contaminants in arctic marine mammals**. *Sci Total Environ* 1994, **154**:107-128.
- Wake K, Motomatsu K, Senoo H, Masuda A, Adachi E: **Improved Kupffer's gold chloride method for demonstrating the stellate cells storing retinol (vitamin A) in the liver and extrahepatic organs of vertebrates**. *Stain Technol* 1986, **61**:193-200.
- Nagy NE, Holven KB, Roos N, Senoo H, Kojima N, Norum KR, Blomhoff R: **Storage of vitamin A in extrahepatic stellate cells in normal rats**. *J Lip Res* 1997, **38**:645-658.
- Brouwer A, Hakansson H, Kukler AK, Van den Berg J, Ahlborg UG: **Marked alterations in retinoid homeostasis of Sprague-Dawley rats induced by a single i.p. dose of 10 micrograms/kg of 2,3,7,8-tetrachlorodibenzo-p-dioxin**. *Toxicology* 1989, **58**:267-283.
- Kelley SK, Nilsson CB, Green MH, Green JB, Hakansson H: **Mobilization of vitamin A stores in rats after administration of 2,3,7,8-tetrachlorodibenzo-p-dioxin: a kinetic analysis**. *Toxicol Sci* 2000, **55**:478-484.
- Skaare JU, Bernhoft A, Derocher A, Gabrielsen GW, Goksøyr A, Henriksen E, Larsen HJ, Lie E, Wiig Ø: **Organochlorines in top predators at Svalbard – occurrence, levels and effects**. *Toxicol Lett* 2000, **112-113**:103-109.