

Proceedings

Open Access

Beta₂-glycoprotein I inhibition of mouse Kupffer cells respiratory burst depends on liver architecture

Ligia F Gomes*¹, Paula R Knox¹, Karin A Simon-Giavarotti²,
Virginia BC Junqueira², Jorge Sans³ and Luis A Videla⁴

Address: ¹Departamento de Análises Clínicas e Toxicológicas, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, São Paulo, Brasil, ²Disciplina de Geriatria, Departamento de Medicina, Universidade Federal de São Paulo, São Paulo, Brasil, ³Unidad de Análisis Celular Integral, ICBM, Facultad de Medicina, Universidad de Chile, Santiago, Chile and ⁴Programa de Farmacología Molecular y Clínica, ICBM, Facultad de Medicina, Universidad de Chile, Santiago, Chile

Email: Ligia F Gomes* - lfgomes@plugnet.com.br; Paula R Knox - paulaknox@hotmail.com; Karin A Simon-Giavarotti - karsimon@bol.com.br; Virginia BC Junqueira - junqueira.dmed@epm.br; Jorge Sans - jsans@machi.med.uchile.cl; Luis A Videla - lvidela@machi.med.uchile.cl

* Corresponding author

from 11th International Symposium on the Cells of the Hepatic Sinusoid and their Relation to Other Cells
Tucson, Arizona, USA, 25–29 August, 2002

Published: 14 January 2004

Comparative Hepatology 2004, **3**(Suppl 1):S43

This article is available from: <http://www.comparative-hepatology.com/content/3/S1/S43>

Introduction

Kupffer cells play important roles in the modulation of immune response, phagocytosis, and senescent cell removal [1,2]. Hydrolytic enzymes and reactive species produce the killing effects of Kupffer cells and some degree of adjacent tissue damage [1,3]. Liver macrophages are constantly exposed to antigens from portal circulation, to which development of full inflammatory response is useless and potentially harmful [4]. Neither tissue damage nor inflammation follows senescent cell removal by Kupffer cells, due to the physiological control of inflammation events during antigen processing [2,5]. Apolipoproteins can modulate macrophage function [6]. Among them, beta₂-glycoprotein I (beta₂GPI) decreases Kupffer cells respiratory burst while increases efficiency of *C. albicans* killing [7]. Beta₂GPI also binds phosphatidylserine (PS) residues on the surface of senescent cells, targeting them to clearance [8]. In order to get an insight on the role of beta₂GPI in the silent antigen removal by Kupffer cells, perfused mouse liver was used as a model of Kupffer cell-dependent phagocytosis and related respiratory burst activity, and results were correlated with those obtained in isolated mouse non-parenchymal cells.

Methods

All reagents used were obtained from Sigma (St. Louis, MO), except for beta₂GPI that was purified from a pool of human sera [7]. Livers from female CF-1 mice (20–28 g

body weight) fed *ad libitum* were perfused with Krebs-Henseleit bicarbonate buffer pH 7.4, saturated with 95% O₂/5% CO₂, at 10 mL/min and 37 degrees C, without recirculation [9]. After 15 min equilibration, O₂ uptake was measured in the effluent perfusate as it flowed past a Clark-type O₂ electrode. Total sinusoidal lactate dehydrogenase (LDH) efflux (in U/g liver) and the respective fractional LDH release (in % of the activity in the tissue) were assessed in the 30–45 min interval as described [9]. Colloidal carbon (C) (0.25 mg/mL; Rotring, Germany) was infused during the 30–45 min interval, either in the absence or presence of 1–30 micrograms beta₂GPI/mL, added at 20 min, and rates of C uptake were calculated according to Cowper et al. [10]. Carbon-induced O₂ consumption (in micromolar O₂/g liver/min) was calculated by subtracting the basal O₂ uptake, during the 30–45 min C perfusion interval [9]. Liver samples taken after perfusion with 0.25 mg C/mL in the absence and in the presence of 30 micrograms beta₂GPI were fixed in Dubosq Brazil, embedded in Paraplast, and stained with hematoxylin-eosin. Non-parenchymal liver cell preparation was obtained by liver perfusion with collagenase [7], with viability values higher than 95%. The respiratory burst was evaluated by a luminol-dependent assay [11] after zymosan stimulation (200 particles/cell) [7], and results were expressed as relative total light emission or light emission rate.





Figure 1
 Infused beta₂-glycoprotein I effects on the perfused mouse liver (A) C-uptake and (B) C-induced O₂ consumption. Means ± SEM for three to five animals/group.




Figure 2
 Structural characteristics of mouse liver parenchyma perfused in vitro with 0.25 mg of colloidal carbon/mL in the (A) absence and (B) presence of 30 micrograms /mL of beta₂-glycoprotein I (beta₂GPI). Haematoxylin-eosin.




Figure 3
 Beta₂-glycoprotein I effects on relative (A) total light emission and (B) light emission rate of isolated cells. Means ± SEM for four separate experiments.

Results and Discussion

Liver perfusion with C in the absence of beta₂GPI led to uptake of C particles and increase in O₂ consumption (Fig. 1, Table 1). The latter effect is mainly related to the respiratory burst of Kupffer cells [4,9], with secondary O₂ utilization in mitochondrial respiration for energy supply needed for C phagocytosis (10) and O₂ uptake induced in hepatocytes by eicosanoids released from activated Kupffer cells [12]. Both Kupffer cell C uptake (Fig. 1A) and C-induced O₂ consumption (Fig. 1B) are inhibited by

Table 1: Carbon (C) uptake and C-induced O₂ uptake inhibition by beta₂-glycoprotein I in perfused mouse liver

Experimental conditions	n	C-uptake (mg/g liver/min)	C-induced O ₂ uptake (μmol/g liver)
Control	5	1.41 ± 0.12	4.82 ± 0.42
Beta ₂ GPI (30 micrograms/mL)	4	1.08 ± 0.005 ^a	0.13 ± 0.13 ^a
Albumin (30 micrograms/mL)	4	1.30 ± 0.06	4.99 ± 0.53

Means ± SEM for separate experiments. ^aP < 0.05 compared to control or albumin infusion, assessed by one-way ANOVA followed by the Bonferroni test.

beta₂GPI (1–30 micrograms/mL), with significant (p < 0.05) 23% and 97% decreases being found at 30 micrograms/L beta₂GPI, respectively. C-induced O₂ consumption inhibition inversely correlates with beta₂GPI concentration (r: -0.8455; p= 0.036). In agreement with biochemical data, optical microscopy revealed that C uptake by non-parenchymal cells is diminished by infusion of 30 micrograms/mL beta₂GPI (Fig. 2). These effects by beta₂GPI are achieved without changes in liver viability, evidenced by comparable fractional LDH effluxes among experimental groups (not shown), and are not mimicked by albumin infusion (Table 1). Despite the beta₂GPI-induced inhibition of C phagocytosis found in perfused liver, chemiluminescence of isolated non-parenchymal liver cells was insensitive to 30 micrograms beta₂GPI/mL (Fig. 3).

Beta₂GPI associates with membranes through annexins, PS receptor, lipoprotein receptors, and negatively charged phospholipids such as PS [5,13]. Interference of beta₂GPI with PS availability in the phagocyte membranes may affect cellular responses, such as translocation of protein kinase C (PKC) to cell membranes [14]. This effect could affect PKC and subsequent triggering of PKC-dependent events, including superoxide anion generation and particle uptake [15]. From current data, it is suggested that beta₂GPI suppresses the respiratory burst response associated with Kupffer cell phagocytosis, while discretely diminishes particle uptake. The former effect is dependent on intact liver architecture, which allows interactions among different cell-types in the liver [16].

Acknowledgements

This work was supported by grants 7000887/1000887 from FONDECYT, Chile (LAV); 97/02335-5 from FAPESP, Brazil (LFG and VBCJ). Fellowships from FAPESP (99/09505-9 to Paula R. Knox/MS; 01/04379-7 to Karin A. Simon-Giavarotti/PD).

References

- Decker K: **The response of liver macrophages to inflammatory stimulation.** *Keio J Med* 1998, **47**:1-9.
- Pradhan D, Krahling S, Williamson P, Schlegel RA: **Multiple systems for recognition of apoptotic lymphocytes by macrophages.** *Mol Biol Cell* 1997, **8**:767-778.
- Wang J-F, Komarov P, de Groot H: **Luminol chemiluminescence in rat macrophages and granulocytes: the role of NO, O₂ / H₂O₂, and HOCl.** *Arch Biochem Biophys* 1993, **304**:189-196.
- Witmer-Pack M, Crowley MT, Inaba K, Steinman RM: **Macrophages, but not dendritic cells, accumulate colloidal carbon following administration in situ.** *J Cell Sci* 1993, **105**:965.
- Fadok VA, Bratton DL, Frasch SC, Warner ML, Henson PM: **The role of phosphatidylserine in recognition of apoptotic cells by phagocytes.** *Cell Death Diff* 1998, **5**:551-562.
- Usynin IF, Tsyrendorzhiev DD, Khar'khovsskii AV, Poliakov LM, Panin LE: **Effect of apolipoprotein A-I on lipopolysaccharide-induced production of reactive oxygen metabolites by Kupffer cells of the rat.** *Biokhimiya, Moscow* 1996, **61**:260-265.
- Gomes LF, Gonçalves LM, Fonseca FLA, Celli CM, Videla LA, Chaimovich H, Junqueira VBC: **Beta₂-Glycoprotein I (apolipoprotein H) modulates uptake and endocytosis associated chemiluminescence in rat Kupffer cells.** *Free Radic Res* 2002, **36**:741-747.
- Chonn A, Semple SC, Cullis PR: **Beta₂-glycoprotein I is a major protein associated with very rapidly cleared liposomes in vivo, suggesting a significant role in the immune clearance of "non-self" particles.** *J Biol Chem* 1995, **270**:25845-25849.
- Videla LA, Tapia G, Fernández V: **Influence of aging on Kupffer cell respiratory activity in relation to particle phagocytosis and oxidative stress parameters in mouse liver.** *Redox Report* 2001, **6**:155-159.
- Cowper KB, Currin RT, Dawson TL, Lindert KA, Lemasters JJ, Thurman RG: **A new method to monitor Kupffer cell function continuously in the perfused rat liver.** *Biochem J* 1990, **266**:141-147.
- Easmon CS, Cole PJ, Williams AJ, Hastings M: **The measurement of opsonic and phagocytic function by luminol chemiluminescence.** *Immunology* 1980, **41**:67-74.
- Qu W, Zhong Z, Goto M, Thurman RG: **Kupffer cell prostaglandin E₂ stimulates parenchymal cell O₂ consumption: alcohol and cell-cell communication.** *Am J Physiol* 1996, **270**:G574-G580.
- Ma K, Simantov R, Zhang J, Silverstein R, Hajjar KA, McCrae KR: **High affinity binding of beta₂-glycoprotein I to human endothelial cells is mediated by annexin II.** *J Biol Chem* 2000, **275**:15541-15548.
- Murray D, Arbuzova A, Hangyás-Mihályiné G, Gambhir A, Ben-Tal N, Honig B, McLaughlin S: **Electrostatic Properties of Membranes Containing Acidic Lipids and Adsorbed Basic Peptides: Theory and Experiment.** *Biophys J* 1999, **77**:3176-3188.
- Dieter P, Schwende H: **Protein kinase C-alpha and -beta play antagonistic roles in the differentiation process of THP-1 cells.** *Cell Signal* 2000, **12**:297-302.
- Potoka DA, Takao S, Owaki T, Bulkley GB, Klein AS: **Endothelial cells potentiate oxidant-mediated Kupffer cell phagocytic killing.** *Free Rad Biol Med* 1998, **24**:1217-1227.